AD		
		_

AWARD NUMBER DAMD17-98-1-8050

TITLE: Cytochrome P450-17alpha Polymorphism and Risk of Breast Cancer

PRINCIPAL INVESTIGATOR: Habibul Ahsan, M.D.

CONTRACTING ORGANIZATION: Columbia University

New York, New York 10032

REPORT DATE: August 1999

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20010109 101

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for the college						UMB No. 0704-0188
Davis Highway, Suite 1204, Arlington, V	/A 22202-430	n is estimated to average 1 hour pating and reviewing the collection ucing this burden, to Washington 1/2, and to the Office of Managem	per response, of information Headquarters ent and Budge	including the time for review Send comments regarding Services, Directorate for Ini t, Paperwork Reduction Pro	wing inst g this bu formation oject (07	tructions, searching existing data sources, riden estimate or any other aspect of this no Operations and Reports, 1215 Jefferson 04-0188), Washington, DC 20503.
I. AGENCY USE ONLY (Leave	e blank)	2. REPORT DATE August 1999	3.	REPORT TYPE AND nnual (1 Aug 98 -	DAT	ES COVERED
4. TITLE AND SUBTITLE Cytochrome P450-17alpha Polymorphism and Risk of Breast Cancer				5. FUNDING NUMBERS DAMD17-98-1-8050		
6. AUTHOR(S) Habibul Ahsan, M.D.			·			
7. PERFORMING ORGANIZATIO Columbia University New York, New York 1003		S) AND ADDRESS(ES)		: :	8. PE	RFORMING ORGANIZATION PORT NUMBER
o. sponsoring / monitoring J.S. Army Medical Research Fort Detrick, Maryland 2170	n and Mat	NAME(S) AND ADDRES	SS(ES)		10. SP	ONSORING / MONITORING SENCY REPORT NUMBER
1. SUPPLEMENTARY NOTES						
2a. DISTRIBUTION / AVAILABI pproved for Public Release;	LITY STAT Distribut	EMENT ion Unlimited		1	2b. D	STRIBUTION CODE
(400 cases and 400 con (LIBSCP). In addition, exposure to exogenous assays on CYP17 polyn Sciences of Columbia U with a digestion by approbeen performed. Delays thus delaying the start of September 1999. Entry questionnaire data (Task preparation and report were stored).	ated with trols) who are a sestrogen norphism. Iniversity in subject of laborate (3) are 6	in an aftered risk of the participated in the also exploring whether are modified by Cons are being conductly. The polymorphis restriction enzymes ext and sample selections analyses. There along data to compute stimated to begin in	breast case Long her the case YP17 potential tend in the case are larger to date to fore, Tasters and n October 1997 and	ancer in a popul Island Breast Careffects of repro- olymorphisms. I he Division of Epeing determine e, assays on 300 the parent LIBS ask 2 will not be merging of labor 1999. Data ar	ation ancer duction the Envirued by Case CP second	ve risk factors and past year, laboratory onmental Health PCR-RFLP methods es and 300 controls have tudy delayed Task 1, apleted until the end of
SUBJECT TERMS cast Cancer, CYP17, estr	ogen, e	epidemiology				15. NUMBER OF PAGES 6 16. PRICE CODE
SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECU OF TH	RITY CLASSIFICATION HIS PAGE Unclassified		JRITY CLASSIFICAT BSTRACT Unclassified	ION	20. LIMITATION OF ABSTRACT

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.
Where copyrighted material is quoted, permission has been obtained to use such material.
Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.
Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.
In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NJY Principalization No. 85-23, Revised 1985).
For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.
In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.
In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.
In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

PI - Signature

Cytochrome P450-17alpha Polymorphism and Risk of Breast Cancer PI: Habibul Ahsan, MD

Table of Contents

Item	Page No.
Form S298	2
Foreword	3
Introduction	5
Body	5-6
Key Research Accomplishments	6
Reportable Outcomes	6
Conclusions	6
References	6

Introduction

Cumulative exposure of circulating estrogen to the breast is linked to the etiology of breast cancer with several studies^{1, 2} suggesting that circulating estradiol (the active form of estrogen) is significantly elevated in breast cancer patients compared to controls in both high and low risk populations³⁻⁶. This variation in circulating estradiol may be due in part to genetic polymorphisms affecting the activity of the enzymes responsible for the metabolism and cellular binding of estrogen in individuals. The enzyme, whose polymorphisms have demonstrated the most potential in the etiology of breast cancer, is the cytochrome P450c17α (CYP17) enzyme. Therefore, the current investigation is investigating whether polymorphisms of the CYP17 gene are associated with an altered risk of breast cancer in a population-based sample of women (400 controls and 400 cases) who participated in the Long Island Breast Cancer Study Project (LIBCSP). Further, we are also exploring whether the effects of reproductive risk factors (e.g., age at menarche, menopause, last child birth and parity) and exposure to exogenous estrogen like hormone replacement therapy, oral contraceptives or estrogen-mimicking substances such as organochlorine pesticides on breast cancer are modified by CYP17 polymorphisms.

Body

Subject and sample selection was delayed in the parent study (LIBSCP) and thus Task I which involved randomly selecting 400 cases and 400 controls (frequency matched to cases by five-year age groups) in batches, using code numbers from among women who had participated in the LIBCSP, was not completed until Month 4. These women provided blood samples and completed a questionnaire for the LIBSCP. Approximately 75% of study participants are postmenopausal in this randomly selected subgroup of the LIBCSP. As expected, the majority of cases and controls randomly selected for this study reflected the racial make-up of the LIBSCP population which is 90% Caucasian, 7% Black and 3% other races.

All laboratory analyses are being done in the laboratory of Dr. Regina Santella in the Division of Environmental Health Sciences of Columbia University. Genomic DNA was isolated from all cases and controls using standard phenol/chloroform extraction method. Polymorphism in the CYP17 gene is being determined by PCR-RFLP methods with the use of appropriate restriction enzymes. Briefly, genomic DNA was amplified by PCR using CYP17-specific primers. Following PCR amplification, digestion of amplified DNA was accomplished by Msp1A1 restriction enzyme. Restriction Fragment Length Polymorphism (RFLP) patterns are being examined by agarose gel electrophoresis after staining with ethidium bromide to detect CYP17 A1 and A2 alleles. The polymorphisms are being identified as follows: 1) a base change in both alleles (A2/A2 genotype) identified if two bands of DNA on the gel are generated, 2) heterozygous (A1/A2) individuals are identified as having three bands on the gel since the enzyme cuts only the mutated allele and not the wild type allele, and 3) individuals who are homzygous (A1/A1) for the alleles produce one band of the uncut PCR product on gel. CYP17 assays completed to date were performed and interpreted blinded to subject characteristics and case-control status. To date these laboratory assays have been performed on only 300 cases and 300 controls as a result of the delay in beginning Task I, as previously mentioned. However, laboratory assays for the remaining cases and controls are expected to be completed by the end of September 1999. Task 3, which includes entry of laboratory data to computers and merging of laboratory data with questionnaire data will be completed in October/November 1999.

Data analysis (Task 4) as well as manuscript preparation and report writing (Task 5) will commence in the coming year.

Key Research Accomplishments

Not applicable at this time.

Reportable Outcomes

Not applicable at this time.

Conclusions

Not applicable at this time.

References:

- 1. Feigelson HS, Henderson BE. Estrogens and breast cancer. Carcinogenesis (Lond) 1996; 17: 2279-84.
- 2. Hulka BS, Liu ET, Lininger RA. Steroid hormones and risk of breast cancer. Cancer 1994: 74(Suppl): 1111-24.
- 3. Bernstein L, Yuan JM, Ross RK, et al. Serum hormone levels in premenopausal Chinese women in Shanghai and white women in Los Angeles: results from two breast cancer case-control studies. Cancer Causes Control 1990; 1:51-8.
- 4. Henderson BE, Gerkins V, Rosario I, Casagrande JT, Pike MC. Elevated serum levels of estrogen and prolactin in daughters of patients with breast cancer. New England J Med 1975;293:790-5.
- 5. Shimizu H, Ross RK, Bernstein L, Pike MC, Henderson BE. Serum estrogen levels in postmenopausal women: comparison of American whites and Japanese in Japan. Br J Cancer 1990; 62: 451-3.
- 6. England P, Skinner L, Cottrell K, Sellwood RA. Serum estradiol 17β in women with benign and malignant breast disease. Br J Cancer 1974; 300:571-6.